Author Index

Aarons, L., see Adedoyin, A. 192
Abouleish, E., see Shao, M.J. 184
Adedoyin, A.
— , Aarons, L. and Houston, J.B.
High-performance liquid chromatographic method for the simultaneous determination of cimetidine and antipyrine in plasma 192
Akamatsu, N., see Ishii, I. 134
Angelo, H.R.
— , Kranz, T., Strøm, J., Thisted, B. and Sørensen, M.B.
High-performance liquid chromatographic method for the determination of dextropropoxyphene and nordextropropoxyphene in serum 413
Aoshima, S., see Oka, K. 419
Baldwin, R.P., see Halbert, M.K. 43
Bargnoux, H., see Bougerolle, A.M. 59
Barreteau, H., see Brouard, A. 453
Battista, H.J.
— , Wehniger, G. and Henn, R.
Separation and identification of nonsteroidal antirheumatic drugs containing a free carboxyl function using high-performance liquid chromatography 77
Bayne, W.F., see Slobodzian, D.K. 345
Belle, S.J.P. Van, see De Smet, M. 309
Benfenati, E., see Farina, P. 323
Benjamin, L.E., see Fell, D. 150
Benson, B., see Gregory, V.M. 140
Berger, J.A., see Bougerolle, A.M. 59
Bougerolle, A.M.
— , Chabard, J.L., Bargnoux, H., Petit, J., Berger, J.A. and Dordain, G.
Simultaneous determination of metapramine and its demethylated metabolites in plasma by gas chromatography—mass spectrometry 59
Brandl, M., see Knoll, R. 425
Brondz, I.
— and Olsen, I.
Sugar composition of lipopolysaccharide from *Haemophilus paraphrophilus* 119
Brooks, M.A.
— and Lee, T.L.
Determination of 4-amino-5-ethyl-3-thiophenecarboxylic acid methyl ester and its acid metabolite in plasma and urine by high-performance liquid chromatography 333
Brosche, T.
— , Platt, D. and Vostrowsky, O.
Methyl enol ethers as artefacts in capillary gas chromatographic profiles of aldehyde dimethyl acetals 219
Brouard, A.
— , Barreteau, H., Merdjan, H., Paillet, M., Micoud, M. and Fredj, G.
Rapid determination of pyrazinamide in biological fluids by high-performance liquid chromatography 453
Brown, N.D.
— , Poon, B.T. and Chulay, J.D.
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 209
Brown, P.R., see Dwyer, M.E. 125
— and Fasching, J.L.
Mobile phase and temperature studies in the reversed-phase liquid chromatography of inosine and guanosine compounds 457
Bullen, W.W., see Gammans, R.E. 285
Caldwell, R.
— and Flanagan, R.J.
Measurement of trazodone in biological fluids by high-performance liquid chromatography 187
Carey, P.F., see Colthup, P.V. 111
Cerulli, N., see Laganà, A. 251
Chabard, J.L., see Bougerolle, A.M. 59
Chavko, M., see Kehr, J. 267
Cheng, L.K., see Osman, M.A. 430
Choi, K.E.
— , Sinkule, J.A., Crom, W.R., Thompson, E.I. and Evans, W.E.
High-performance liquid chromatographic assay of mitomycin in biological fluids 197
Chulay, J.D., see Brown, N.D. 209
Churchill, F.C., see Dickinson, C.M. 91
Cohen, A.I., see Jemal, M. 299
Colthup, P.V.
— , Dallas, F.A.A., Saynor, D.A., Carey, P.F., Skidmore, I.F. and Martin, L.E.
Determination of salbutamol in human plasma and urine by high-performance thin-layer chromatography 111
Crom, W.R., see Choi, K.E. 197
Dallas, F.A.A., see Colthup, P.V. 111
De Deyn, P., see Marescau, B. 215
De Smet, M.
— , Van Belle, S.J.P., Storme, G.A. and Massart, D.L.
High-performance liquid chromato-

Author Index

Aarons, L., see Adedoyin, A. 192
Abouleish, E., see Shao, M.J. 184
Adedoyin, A.
— , Aarons, L. and Houston, J.B.
High-performance liquid chromatographic method for the simultaneous determination of cimetidine and antipyrine in plasma 192
Akamatsu, N., see Ishii, I. 134
Angelo, H.R.
— , Kranz, T., Strøm, J., Thisted, B. and Sørensen, M.B.
High-performance liquid chromatographic method for the determination of dextropropoxyphene and nordextropropoxyphene in serum 413
Aoshima, S., see Oka, K. 419
Baldwin, R.P., see Halbert, M.K. 43
Bargnoux, H., see Bougerolle, A.M. 59
Barreteau, H., see Brouard, A. 453
Battista, H.J.
— , Wehniger, G. and Henn, R.
Separation and identification of nonsteroidal antirheumatic drugs containing a free carboxyl function using high-performance liquid chromatography 77
Bayne, W.F., see Slobodzian, D.K. 345
Belle, S.J.P. Van, see De Smet, M. 309
Benfenati, E., see Farina, P. 323
Benjamin, L.E., see Fell, D. 150
Benson, B., see Gregory, V.M. 140
Berger, J.A., see Bougerolle, A.M. 59
Bougerolle, A.M.
— , Chabard, J.L., Bargnoux, H., Petit, J., Berger, J.A. and Dordain, G.
Simultaneous determination of metapramine and its demethylated metabolites in plasma by gas chromatography—mass spectrometry 59
Brandl, M., see Knoll, R. 425
Brondz, I.
— and Olsen, I.
Sugar composition of lipopolysaccharide from *Haemophilus paraphrophilus* 119
Brooks, M.A.
— and Lee, T.L.
Determination of 4-amino-5-ethyl-3-thiophenecarboxylic acid methyl ester and its acid metabolite in plasma and urine by high-performance liquid chromatography 333
Brosche, T.
— , Platt, D. and Vostrowsky, O.
Methyl enol ethers as artefacts in capillary gas chromatographic profiles of aldehyde dimethyl acetals 219
Brouard, A.
— , Barreteau, H., Merdjan, H., Paillet, M., Micoud, M. and Fredj, G.
Rapid determination of pyrazinamide in biological fluids by high-performance liquid chromatography 453
Brown, N.D.
— , Poon, B.T. and Chulay, J.D.
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 209
Brown, P.R., see Dwyer, M.E. 125
— and Fasching, J.L.
Mobile phase and temperature studies in the reversed-phase liquid chromatography of inosine and guanosine compounds 457
Bullen, W.W., see Gammans, R.E. 285
Caldwell, R.
— and Flanagan, R.J.
Measurement of trazodone in biological fluids by high-performance liquid chromatography 187
Carey, P.F., see Colthup, P.V. 111
Cerulli, N., see Laganà, A. 251
Chabard, J.L., see Bougerolle, A.M. 59
Chavko, M., see Kehr, J. 267
Cheng, L.K., see Osman, M.A. 430
Choi, K.E.
— , Sinkule, J.A., Crom, W.R., Thompson, E.I. and Evans, W.E.
High-performance liquid chromatographic assay of mitomycin in biological fluids 197
Chulay, J.D., see Brown, N.D. 209
Churchill, F.C., see Dickinson, C.M. 91
Cohen, A.I., see Jemal, M. 299
Colthup, P.V.
— , Dallas, F.A.A., Saynor, D.A., Carey, P.F., Skidmore, I.F. and Martin, L.E.
Determination of salbutamol in human plasma and urine by high-performance thin-layer chromatography 111
Crom, W.R., see Choi, K.E. 197
Dallas, F.A.A., see Colthup, P.V. 111
De Deyn, P., see Marescau, B. 215
De Smet, M.
— , Van Belle, S.J.P., Storme, G.A. and Massart, D.L.
High-performance liquid chromato-

graphic determination of vinca-alkaloids in plasma and urine 309
Deyn, P. De, see Marescau, B. 215
Dickinson, C.M.
— , Navin, T.R. and Churchill, F.C.
High-performance liquid chromatographic method for quantitation of pentamidine in blood serum 91
Dikkeschei, L.D.
— , Van Veelen, H., Nagel, G.T., Willemse, P.H.B. and Wolthers, B.G.
Specific and sensitive determination of medroxyprogesterone acetate in human serum by gas chromatography—mass spectrometry 1
D'Incalci, M., see Farina, P. 323
Dordain, G., see Bougerolle, A.M. 59
Duval, M., see Godbillon, J. 365
Dwyer, M.E.
— and Brown, P.R.
Evaluation of combined electrochemical and ultraviolet detection in the high-performance liquid chromatographic analysis of plasma for low-molecular weight constituents 125
Ebihara, M., see Goto, J. 241
Ebinger, G., see Herregodts, P. 33
Evans, W.E., see Choi, K.E. 197
Fallon, K.D., see Shao, M.J. 184
Farina, P.
— , Benfenati, E., Lassiani, L., Nisi, C. and D'Incalci, M.
High-performance liquid chromatographic assay for the determination of *p*-(3,3-dimethyl-1-triazeno)benzoic acid in mouse plasma 323
Fasching, J.L., see Brown, P.R. 457
Fell, D.
— , Benjamin, L.E. and Steele, R.D.
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by high-performance liquid chromatography 150
Flanagan, R.J., see Calwell, R. 187
Fredj, G., see Brouard, A. 453
Fujiwara, T., see Tokuma, Y. 51
Galushko, S.V.
— and Shishkina, I.P.
Reversed-phase ion-paired high-performance liquid chromatographic determination of 6-azacytidine in blood 157
Gammans, R.E.
— , Kerns, E.H. and Bullen, W.W.
Capillary gas chromatographic—mass spectrometric determination of buspirone in plasma 285
Gifford, R., see Swagzdis, J.E. 203
Godbillon, J.
— , Duval, M. and Gosset, G.
Determination of oxprenolol in human plasma by high-performance liquid chromatography, in comparison with gas chromatography and gas chromatography—mass spectrometry 365
Goebel, K.-J.
— and Kölle, E.U.
High-performance liquid chromatographic determination of diltiazem and four of its metabolites in plasma. Application to pharmacokinetics 355
Gorp, L. Van, see Marescau, B. 215
Gosset, G., see Godbillon, J. 365
Goto, J.
— , Suzaki, K., Ebihara, M., Nambara, T. and Masu, A.
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glururonides in human urine by high-performance liquid chromatography 241
Goulle, J.P., see Lacroix, C. 436
Gregory, V.M.
— , Larsen, B. and Benson, B.
Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by high-performance liquid chromatography using electrochemical detection 140
Gringore, A., see Lacroix, C. 436
Hagen, B.
— , Walseth, F., Walstad, R.A. and Iversen, T.
Gas chromatographic assay of triethylenethiophosphoramide in serum and urine 173
Halbert, M.K.
— and Baldwin, R.P.
Determination of cysteine and glutathione in plasma and blood by liquid chromatography with electrochemical detection using a chemically modified electrode containing cobalt phthalocyanine 43
Hayashi, R., see Nagashima, H. 373
Heinemann, V., see Schwandt, P. 145
Henn, R., see Battista, H.J. 77
Henning, H.V., see Schneider, K. 19
Herregodts, P.
— , Michotte, Y. and Ebinger, G.
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a com-

bined extraction procedure and high-performance liquid chromatography with electrochemical detection 33
Holland, M.L.
— and Ng, K.T.
Automated capillary gas chromatographic assay using nitrogen—phosphorus detection for the determination of cetiedil in plasma 178
Horai, Y.
— and Ishizaki, T.
Rapid and sensitive liquid chromatographic method for the determination of dapsone and monoacetyldapsone in plasma and urine 447
Hosoi, M., see Maeda, K. 11
Houston, J.B., see Adedoyin, A. 192
Hsieh, J.Y.-K., see Slobodzian, D.K. 345
Ishida, J., see Kai, M. 259
Ishii, I.
— , Takahashi, N., Kato, S., Akamatsu, N. and Kawazoe, Y.
High-performance liquid chromatographic analysis of changes of asparagine-linked oligosaccharides in regenerating rat liver 134
Ishizaki, T., see Horai, Y. 447
Ivashkiv, E., see Jemal, M. 299
Iversen, T., see Hagen, B. 173
Jemal, M.
— , Ivashkiv, E., Ribick, M. and Cohen, A.I.
Determination of SQ 27,519, the active phosphinic acid—carboxylic acid of the prodrug SQ 28,555, in human serum by capillary gas chromatography with nitrogen—phosphorus detection after a two-step derivatization 299
Kai, M.
— , Miura, T., Ishida, J. and Ohkura, Y.
High-performance liquid chromatographic method for monitoring leupeptin in mouse serum and muscle by pre-column fluorescence derivatization with benzoin 259
Kato, S., see Ishii, I. 134
Kawazoe, Y., see Ishii, I. 134
Kehr, J.
— and Chavko, M.
Rapid determination of adenine nucleotides in brain tissue by ion-paired reversed-phase column liquid chromatography under isocratic conditions 267
Kerns, E.H., see Gammans, R.E. 285
Khalil, S.N., see Shao, M.J. 184
Knoll, R.
— and Brandl, M.
Rapid and simple method for the routine determination of dobutamine in human plasma by high-performance liquid chromatography 425
Kölle, E.U., see Goebel, K.-J. 355
Kranz, T., see Angelo, H.R. 413
Kroschel, D.M., see Powis, G. 402
Lacroix, C.
— , Levert, P., Laine, G., Goulle, J.P. and Gringore, A.
Microdosage de la 5-fluorocytosine par chromatographie en phase liquide et détection fluorimétrique 436
Laganà, A.
— , Liberti, A., Politi, L., Scandurra, R. and Cerulli, N.
Chromatographic evaluation of perfusion on charcoal in ureamia 251
Laine, G., see Lacroix, C. 436
Larsen, B., see Gregory, V.M. 140
Lassiani, L., see Farina, P. 323
Lee, T.L., see Brooks, M.A. 333
Lennard, L.
Assay of 6-mercaptopurine in human plasma 441
Lester, D.C., see Nierenberg, D.W. 275
Levert, P., see Lacroix, C. 436
Liberti, A., see Laganà, A. 251
Lin, K.-T.
— , Momparler, R.L. and Rivard, G.E.
Sample preparation for the determination of 5-aza-2'deoxycytidine in plasma by high-performance liquid chromatography 162
Lowenthal, A., see Marescau, B. 215
Lu, K.
— , Savaraj, N. and Newman, R.A.
High-performance liquid chromatographic separation and quantitation of benzisoquinolinedione (Nafidimide) from biological fluids 408
Maeda, K.
— , Shiraishi, S., Sakamoto, N., Ohki, T., Hosoi, M., Ohta, K. and Yamanaka, N.
Identification of *Escherichia coli* by detection of hydroquinone and uracil in the urine system 11
Marescau, B.
— , De Deyn, P., Wiechert, P., Van Gorp, L. and Lowenthal, A.
Quantitative overestimation of free taurocyamine using short liquid cation-exchange chromatography columns? 215
Martin, L.E., see Colthup, P.V. 111
Marzo, A.
— and Treffner, E.
Gas—liquid chromatographic determina-

tion of isosorbide-5-mononitrate derivatized with perfluorinated anhydrides for pharmacokinetic and bioavailability investigations 390
Massart, D.L., see De Smet, M. 309
Masu, A., see Goto, J. 241
Matthaei, D., see Schneider, K. 19
Merdjan, H., see Brouard, A. 453
Michotte, Y., see Herregodts, P. 33
Mico, B.A., see Swagzdis, J.E. 203
Micoud, M., see Brouard, A. 453
Miura, T., see Kai, M. 259
Momparler, R.L., see Lin, K.-T. 162
Moore, D.J., see Powis, G. 402
Morreal, C.E.
— , Sinha, D.K., White, C.J. and Nemoto, D.T.
Assay of prostaglandins in the epithelial cells and fibroblasts of the rat mammary gland 380
Mutschler, E., see Schladitz-Keil, G. 99
Nagashima, H.
— , Tanaka, Y. and Hayashi, R.
Column liquid chromatography for the simultaneous determination of the enantiomers of loxoprofen sodium and its metabolites in human urine 373
Nagel, G.T., see Dikkeschei, L.D. 1
Nambara, T., see Goto, J. 241
Navin, T.R., see Dickinson, C.M. 91
Nemoto, D.T., see Morreal, C.E. 380
Neupert, M., see Schneider, K. 19
Newman, R.A., see Lu, K. 408
Ng, K.T., see Holland, M.L. 178
Nierenberg, D.W.
— and Lester, D.C.
Determination of vitamins A and E in serum and plasma using a simplified clarification method and high-performance liquid chromatography 275
Nisi, C., see Farina, P. 323
Noguchi, M., see Oka, K. 419
— , see Tokuma, Y. 51
O'Hare, M.J., see Smith, J.A. 168
Ohki, T., see Maeda, K. 11
Ohkura, Y., see Kai, M. 259
Ohta, K., see Maeda, K. 11
Oka, K.
— , Aoshima, S. and Noguchi, M.
Highly sensitive determination of ketoprofen in human serum and urine and its application to pharmacokinetic study 419
Olsen, I., see Brondz, I. 119
Osman, M.A.
— , Pinchbeck, F.M., Cheng, L.K. and Wright, G.J.
Determination of lenperone in dog plasma by high-performance liquid chromatography 430
Paar, W.D., see Von Unruh, G.E. 67
Paillet, M., see Brouard, A. 453
Petit, J., see Bougerolle, A.M. 59
Pfäffli, P.
— and Svartling, N.
Assay of methyl methacrylate in blood samples by headspace capillary gas chromatography 386
Pinchbeck, F.M., see Osman, M.A. 430
Platt, D., see Brosche, T. 219
Politi, L., see Laganà, A. 251
Poon, B.T., see Brown, N.D. 209
Powis, G.
— , Kroschel, D.M. and Moore, D.J.
Sensitive reversed-phase high performance liquid chromatographic assay for the antitumor agent benzisoquinolinedione (Nafidimide) employing direct on-column injection of plasma and urine, and its use for pharmacokinetic studies in a dog 402
Rakhit, A., see Tipnis, V. 396
Ribick, M., see Jemal, M. 299
Richter, W.O., see Schwandt, P. 145
Rivard, G.E., see Lin, K.-T. 162
Sakamoto, N., see Maeda, K. 11
Savaraj, N., see Lu, K. 408
Saynor, D.A., see Colthup, P.V. 111
Scandurra, R., see Laganà, A. 251
Scheler, F., see Schneider, K. 19
Schladitz-Keil, G.
— , Spahn, H. and Mutschler, E.
Fluorimetric determination of the quaternary compound trospium and its metabolite in biological material after derivatization with benoxaprofen chloride 99
Schneider, K.
— , Neupert, M., Spiteller, G., Henning, H.V., Matthaei, D. and Scheler, F.
Gas chromatography of amino acids in urine and haemofiltrate 19
Schwandt, P.
— , Richter, W.O., Heinemann, V. and Weisweiler, P.
Characterization of human apolipoprotein A-I by reversed-phase high-performance liquid chromatography 145
Shao, M.J.
— , Fallon, K.D., Khalil, S.N. and Abouleish, E.
Quantitation of gallamine (Flaxedil) in human plasma using high-performance liquid chromatography 184

Shiraishi, S., see Maeda, K. 11
Shishkina, I.P., see Galushko, S.V. 157
Sinha, D.K., see Morreal, C.E. 380
Sinkule, J.A., see Choi, K.E. 197
Skidmore, I.F., see Colthup, P.V. 111
Slobodzian, D.K.
— , Hsieh, J.Y.-K. and Bayne, W.F.
Simultaneous determination of tranilast and metabolites in plasma and urine using high-performance liquid chromatography 345
Smet, M. De, see De Smet, M. 309
Smith, J.A.
— and O'Hare, M.J.
Ion-exchange high-performance liquid chromatography of mouse epidermal growth factor and its congeners: mobile phase optimization with ion-pairing additives 168
Sørensen, M.B., see Angelo, H.R. 413
Spahn, H., see Schladitz-Keil, G. 99
Spiteller, G., see Schneider, K. 19
Steele, R.D., see Fell, D. 150
Storme, G.A., see De Smet, M. 309
Strøm, J., see Angelo, H.R. 413
Suzaki, K., see Goto, J. 241
Svartling, N., see Pfäffli, P. 386
Swagzdis, J.E.
— , Gifford, R. and Mico, B.A.
Determination of 4-(2-di-*n*-propylaminoethyl)-7-hydroxy-2-(3H)-indolone and N-[2′-hydroxy-5′-(N,N-di-*n*-propylaminoethylphenyl)]methane sulfonamide in plasma by high-performance liquid chromatography with electrochemical detection 203
Takahashi, N., see Ishii, I. 134
Tanaka, Y., see Nagashima, H. 373
Thisted, B., see Angelo, H.R. 413
Thompson, E.I., see Choi, K.E. 197
Tipnis, V.
— and Rakhit, A.
Determination of pentopril, an angiotension converting enzyme inhibitor, and its active metabolite in urine 396
Tokuma, Y.
— , Fujiwara, T. and Noguchi, H.
Determination of nilvadipine in human plasma by capillary column gas chromatography—negative-ion chemical-ionization mass spectrometry 51
Treffner, E., see Marzo, A. 390
Unruh, G.E. Von, see Von Unruh, G.E. 67
Urayama, T., see Yanagisawa, I. 229
Van Belle, S.J.P., see De Smet, M. 309
Van Gorp, L., see Marescau, B. 215
Van Veelen, H., see Dikkeschei, L.D. 1
Veelen, H. van, see Dikkeschei, L.D. 1
Von Unruh, G.E.
— and Paar, W.D.
Gas chromatographic assay for oxcarbazepine and its main metabolites in plasma 67
Vostrowsky, O., see Brosche, T. 219
Walseth, F., see Hagen, B. 173
Walstad, R.A., see Hagen, B. 173
Wehninger, G., see Battista, H.J. 77
Weisweiler, P., see Schwandt, P. 145
White, C.J., see Morreal, C.E. 380
Wiechert, P., see Marescau, B. 215
Willemse, P.H.B., see Dikkeschei, L.D. 1
Wilson, K., see Colthup, P.V. 111
Wolthers, B.G., see Dikkeschei, L.D. 1
Wright, G.J., see Osman, M.A. 430
Yamanaka, N., see Maeda, K. 11
Yamane, M., see Yanagisawa, I. 229
Yanagisawa, I.
— , Yamane, M. and Urayama, T.
Simultaneous separation and sensitive determination of free fatty acids in blood plasma by high-performance liquid chromatography 229

Subject Index

Adenine nucleotides
Rapid determination of adenine nucleotides in brain tissue by ion-paired reversed-phase column LC under isocratic conditions 267

Adenosine
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 150

S-Adenosylethionine
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 150

S-Adenosylhomocysteine
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 150

S-Adenosylmethionine
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 150

ADP
Rapid determination of adenine nucleotides in brain tissue by ion-paired reversed-phase column LC under isocratic conditions 267

Aldehyde dimethyl acetals
Methyl enol ethers as artefacts in capillary GC profiles of aldehyde dimethyl acetals 219

Amines
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33

Amino acids
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 150

Amino acids
GC of amino acids in urine and haemofiltrate 19

4-Amino-5-ethyi-3-thiophenecarboxylic acid
Determination of 4-amino-5-ethyl-3-thiophenecarboxylic acid methyl ester and its acid metabolite in plasma and urine by HPLC 333

4-Amino-5-ethyl-3-thiophenecarboxylic acid methyl ester
Determination of 4-amino-5-ethyl-3-thiophenecarboxylic acid methyl ester and its acid metabolite in plasma and urine by HPLC 333

AMP
Rapid determination of adenine nucleotides in brain tissue by ion-paired reversed-phase column LC under isocratic conditions 267

Antipyrine
HPLC method for the simultaneous determination of cimetidine and antipyrine in plasma 192

Apolipoprotein A-I
Characterization of human apolipoprotein A-I by reversed-phase HPLC 145

Asparagine-linked oligosaccharides
HPLC analysis of changes of asparagine-linked oligosaccharides in regenerating rat liver 134

ATP
Rapid determination of adenine nucleotides in brain tissue by ion-paired reversed-phase column LC under isocratic conditions 267

6-Azacytidine
Reversed-phase ion-paired HPLC determination of 6-azacytidine in blood 157

5-Aza-2′-deoxycytidine
Sample preparation for the determination of 5-aza-2′-deoxycytidine in plasma by HPLC 162

Benzisoquinolinedione
HPLC separation and quantitation of benzisoquinolinedione (Nafidimide) from biological fluids 408

Benzisoquinolinedione
Sensitive reversed-phase HPLC assay for the antitumor agent benzisoquinolinedione (Nafidimide) employing direct on-column injection of plasma and urine, and its use for pharmacokinetic studies in a dog 402

Bile acid 3-glucuronides
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 241

Biogenic amines
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33

Buspirone
Capillary GC—MS determination of buspirone in plasma 285

Catecholamines
Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 140
Cetiedil
Automated capillary GC assay using nitrogen—phosphorus detection for the determination of cetiedil in plasma 178
Charcoal
Chromatographic evaluation of perfusion on charcoal in uraemia 251
Chenodeoxycholate 3-glucuronide
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 241
7-Chloro-4-aminoquinoline
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 209
7-Chloro-4-hydroxyquinoline
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 209
Chloroquine
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 209
Cholate 3-glucuronide
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 241
Cimetidine
HPLC method for the simultaneous determination of cimetidine and antipyrine in plasma 192
Cysteine
Determination of cysteine and glutathione in plasma and blood by LC with electrochemical detection using a chemically modified electrode containing cobalt phthalocyanine 43
Dapsone
Rapid and sensitive LC method for the determination of dapsone and monoacetyldapsone in plasma and urine 447
Deoxycholate 3-glucuronide
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 241
Dextropropoxyphene
HPLC method for the determination of dextropropoxyphene and nordextropropoxyphene in serum 413
trans-10,11-Dihydroxycarbazepine
GC assay for oxcarbazepine and its main metabolites in plasma 67
3,4-Dihydroxyphenylacetic acid
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33
Diltiazem
HPLC determination of diltiazem and four of its metabolites in plasma. Application to pharmacokinetics 355
N-(3′,4′-Dimethoxycinnamoyl)anthranilic acid
Simultaneous determination of tranilast and metabolites in plasma and urine using HPLC 345
Dimethyl acetals
Methyl enol ethers as artefacts in capillary GC profiles of aldehyde dimethyl acetals 219
p-(3,3-Dimethyl-1-triazeno)benzoic acid
HPLC assay for the determination of *p*-(3,3-dimethyl-1-triazeno)benzoic acid in mouse plasma 323
4-(2-Di-*n*-propylaminoethyl)-7-hydroxy-2-(3H)-indolone
Determination of 4-(2-di-*n*-propylaminoethyl)-7-hydroxy-2-(3H)-indolone and N-[2′-hydroxy-5′-(N,N-di-*n*-propylaminoethylphenyl)]methane sulfonamide in plasma by HPLC with electrochemical detection 203
Dobutamine
Rapid and simple method for the routine determination of dobutamine in human plasma by HPLC 425
Dopamine
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33
Dopamine
Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 140
Drugs
Separation and identification of nonsteroidal antirheumatic drugs containing a free carboxyl function using HPLC 77

Epidermal growth factor
Ion-exchange HPLC of mouse epidermal growth factor and its congeners: mobile phase optimization with ion-pairing additives 168
Epinephrine
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33
Escherichia coli
Identification of *Escherichia coli* by detection of hydroquinone and uracil in the urine system 11
Fatty acids
Simultaneous separation and sensitive determination of free fatty acids in blood plasma by HPLC 229
Fenac compounds
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 77
Fenbufen
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 77
Fenemates
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 77
Flaxedil
Quantitation of gallamine (Flaxedil) in human plasma using HPLC 184
5-Fluorocystosine
Determination of 5-fluorocytosine by HPLC with fluorometric detection 436
Free fatty acids
Simultaneous separation and sensitive determination of free fatty acids in blood plasma by HPLC 229
Gallamine
Quantitation of gallamine (Flaxedil) in human plasma using HPLC 184
Glutathione
Determination of cysteine and glutathione in plasma and blood by LC with electrochemical detection using a chemically modified electrode containing cobalt phthalocyanine 43
Glycochenodeoxycholate 3-glucuronide
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 241
Glycocholate 3-glucuronide
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 241
Glycoproteins
HPLC analysis of changes of asparagine-linked oligosaccharides in regenerating rat liver 134
Guanidino compounds
Quantitative overestimation of free taurocyamine using short liquid cation-exchange chomatography columns? 215
Guanosine
Evaluation of combined electrochemical and ultraviolet detection in the HPLC analysis of plasma for low-molecular-weight constituents 125
Guanosine
Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 457
Guanosine 5'-phosphate
Evaluation of combined electrochemical and ultraviolet detection in the HPLC analysis of plasma for low-molecular-weight constituents 125
Haemophilus paraphrophilus
Sugar composition of lipopolysaccharide from *Haemophilus paraphrophilus* 119
High-density lipoproteins
Characterization of human apolipoprotein A-I by reversed-phase HPLC 145
Homovanillic acid
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33
Hydroquinone
Identification of *Escherichia coli* by detection of hydroquinone and uracil in the urine system 11
10-Hydroxycarbazepine
GC assay for oxcarbazepine and its main metabolites in plasma 67
5-Hydroxyindoleacetic acid
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33

5-Hydroxyindoleacetic acid
Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 140
4-(4'-Hydroxy-1'-methylbutylamino)-7-chloroquinoline
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 209
N-[2'-Hydroxy-5'-(N,N-di-*n*-propylaminoethylphenyl)]methane sulphonamide
Determination of 4-(2-di-*n*-propylaminoethyl)-7-hydroxy-2-(3H)-indolone and N-[2'-hydroxy-5'-(N,N-di-*n*-propylaminoethylphenyl)]methane sulfonamide in plasma by HPLC with electrochemical detection 203
5-Hydroxytryptamine
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33
5-Hydroxytryptamine
Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 140
Hypoxanthine
Evaluation of combined electrochemical and ultraviolet detection in the HPLC analysis of plasma for low-molecular-weight constituents 125
Inosine
Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 457
Isosorbide-5-mononitrate
GLC determination of isosorbide-5-mononitrate derivatized with perfluorinated anhydrides for pharmacokinetic and bioavailability investigations 390
Ketoprofen
Highly sensitive determination of ketoprofen in human serum and urine and its application to pharmacokinetic study 419
Lenperone
Determination of lenperone in dog plasma by HPLC 430
Leupeptin
HPLC method for monitoring leupeptin in mouse serum and muscle by pre-column fluorescence derivatization with benzoin 259
Lipopolysaccharides
Sugar composition of lipopolysaccharide from *Haemophilus paraphrophilus* 119
Lipoproteins
Characterization of human apolipoprotein A-I by reversed-phase HPLC 145
Loxoprofen
Column LC for the simultaneous determination of the enantiomers of loxoprofen sodium and its metabolites in human urine 373
Loxoprofen, hydroxy metabolites
Column LC for the simultaneous determination of the enantiomers of loxoprofen sodium and its metabolites in human urine 373
Medroxyprogesterone acetate
Specific and sensitive determination of medroxyprogesterone acetate in human serum by GC—MS 1
6-Mercaptopurine
Assay of 6-mercaptopurine in human plasma 441
Metapramine
Simultaneous determination of metapramine and its demethylated metabolites in plasma by GC—MS 59
3-Methoxy-4-hydroxyphenylethylene glycol
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33
3-Methoxytyramine
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33
Methyl enol ethers
Methyl enol ethers as artefacts in capillary GC profiles of aldehyde dimethyl acetals 219
Methyl methacrylate
Assay of methyl methacrylate in blood samples by headspace capillary GC 386
Mitomycin
HPLC assay of mitomycin in biological fluids 197
Monoacetyldapsone
Rapid and sensitive LC method for the determination of dapsone and monoacetyldapsone in plasma and urine 447

Nafidimide
HPLC separation and quantitation of benzisoquinolinedione (Nafidimide) from biological fluids 408
Nafidimide
Sensitive reversed-phase HPLC assay for the antitumor agent benzisoquinolinedione (Nafidimide) employing direct on-column injection of plasma and urine, and its use for pharmacokinetic studies in a dog 402
Nilvadipine
Determination of nilvadipine in human plasma by capillary column GC—negative-ion chemical-ionization MS 51
Non-steroidal anti-inflammatory drugs
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 77
Nordextropropoxyphene
HPLC method for the determination of dextropropoxyphene and nordextropropoxyphene in serum 413
Norepinephrine
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33
Norepinephrine
Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 140
Oligosaccharides
HPLC analysis of changes of asparagine-linked oligosaccharides in regenerating rat liver 134
Oxcarbazepine
GC assay for oxcarbazepine and its main metabolites in plasma 67
Oxprenolol
Determination of oxprenolol in human plasma by HPLC, in comparison with GC and GC—MS 365
Pentamidine
HPLC method for quantitation of pentamidine in blood serum 91
Pentopril
Determination of pentopril, an angiotensin converting enzyme inhibitor, and its active metabolite in urine 396
Profens
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 77
Prostaglandins
Assay of prostaglandins in the epithelial cells and fibroblasts of the rat mammary gland 380
Pyrazinamide
Rapid determination of pyrazinamide in biological fluids by HPLC 453
Retinol
Determination of vitamins A and E in serum and plasma using a simplified clarification method and HPLC 275
Salbutamol
Determination of salbutamol in human plasma and urine by HPTLC 111
Salicylates
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 77
Serotonin
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 33
SQ 27,519
Determination of SQ 27,519, the active phosphinic acid—carboxylic acid of the prodrug SQ 28,555, in human serum by capillary GC with nitrogen—phosphorus detection after a two-step derivatization 299
SQ 28,555
Determination of SQ 27,519, the active phosphinic acid—carboxylic acid of the prodrug SQ 28,555, in human serum by capillary GC with nitrogen—phosphorus detection after a two-step derivatization 299
Steroids
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 241
Sugars
Sugar composition of lipopolysaccharide from *Haemophilus paraphrophilus* 119
Sulphur amino acids
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 150
Taurochenodeoxycholate 3-glucuronide
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 241
Taurocyamine
Quantitative overestimation of free taurocyamine using short liquid cation-exchange chromatography columns? 215

α-Tocopherol
Determination of vitamins A and E in serum and plasma using a simplified clarification method and HPLC 275
Tranilast
Simultaneous determination of tranilast and metabolites in plasma and urine using HPLC 345
Trazodone
Measurement of trazodone in biological fluids by HPLC 187
Triethylenethiophosphoramide
GC assay of triethylenethiophosphoramide in serum and urine 173
Trospium
Fluorimetric determination of the quaternary compound trospium and its metabolite in biological material after derivatization with benoxaprofen chloride 99
L-Tryptophan
Evaluation of combined electrochemical and ultraviolet detection in the HPLC analysis of plasma for low-molecular-weight constituents 125
Tyrosine
Evaluation of combined electrochemical and ultraviolet detection in the HPLC analysis of plasma for low-molecular-weight constituents 125
Uracil
Identification of *Escherichia coli* by detection of hydroquinone and uracil in the urine system 11
Uric acid
Evaluation of combined electrochemical and ultraviolet detection in the HPLC analysis of plasma for low-molecular-weight constituents 125
Vinblastine
HPLC determination of vinca-alkaloids in plasma and urine 309
Vinca-alkaloids
HPLC determination of vinca-alkaloids in plasma and urine 309
Vincristine
HPLC determination of vinca-alkaloids in plasma and urine 309
Vindesine
HPLC determination of vinca-alkaloids in plasma and urine 309
Vitamins
Determination of vitamins A and E in serum and plasma using a simplified clarification method and HPLC 275
Xanthine
Evaluation of combined electrochemical and ultraviolet detection in the HPLC analysis of plasma for low-molecular-weight constituents 125
Xanthosine 5′-phosphate
Evaluation of combined electrochemical and ultraviolet detection in the HPLC analysis of plasma for low-molecular-weight constituents 125